

## Bayesian networks in healthcare: What is preventing their adoption?

Evangelia Kyrimi<sup>a,\*</sup>, Kudakwashe Dube<sup>b,c</sup>, Norman Fenton<sup>a</sup>, Ali Fahmi<sup>a</sup>,  
Mariana Ranieri Neves<sup>a</sup>, William Marsh<sup>a</sup>, Scott McLachlan<sup>a,b</sup>

<sup>a</sup> School of Electronic Engineering & Computer Science, Queen Mary University of London, Mile End Road, London, E1 4NS, UK

<sup>b</sup> Health Informatics and Knowledge Engineering Research (HiKER) Group

<sup>c</sup> School of Fundamental Sciences, Massey University, Palmerston North, 4442, New Zealand

### ARTICLE INFO

#### Keywords:

Bayesian networks  
Adoption  
Implementation  
Healthcare

### ABSTRACT

There has been much research effort expended toward the use of Bayesian networks (BNs) in medical decision-making. However, because of the gap between developing an accurate BN and demonstrating its clinical usefulness, this has not resulted in any widespread BN adoption in clinical practice. This paper investigates this problem with the aim of finding an explanation and ways to address the problem through a comprehensive literature review of articles describing BNs in healthcare. Based on the literature collection that has been systematically narrowed down from 3810 to 116 most relevant articles, this paper analyses the *benefits*, *barriers* and *facilitating factors* (BBF) for implementing BN-based systems in healthcare using the ITPOSMO-BBF framework. A key finding is that works in the literature rarely consider *barriers* and even when these were identified they were not connected to *facilitating factors*. The main finding is that the barriers can be grouped into: (1) data inadequacies; (2) clinicians' resistance to new technologies; (3) lack of clinical credibility; (4) failure to demonstrate clinical impact; (5) absence of an acceptable predictive performance; and (6) absence of evidence for model's generalisability. The facilitating factors can be grouped into: (1) data collection improvements; (2) software and technological improvements; (3) having interpretable and easy to use BN-based systems; (4) clinical involvement in the development or review of the model; (5) investigation of model's clinical impact; (6) internal validation of the model's performance; and (7) external validation of the model. These groupings form a strong basis for a generic framework that could be used for formulating strategies for ensuring BN-based clinical decision-support system adoption in frontline care settings. The output of this review is expected to enhance the dialogue among researchers by providing a deeper understanding for the neglected issue of BN adoption in practice and promoting efforts for implementing BN-based systems.

### 1. Introduction

In healthcare, clinicians make countless decisions in daily practice. These decisions include estimating the likelihood that the patient is suffering from a particular disease (diagnostic decisions), that the chosen intervention represents the optimal treatment strategy for the patient's condition (treatment decisions), or that the selected treatment will result in a certain outcome (prognostic decisions). The vast array of information now available, as well as the complex nature of human physiology, all serve to increase uncertainty and make clinical decision-making a challenging task [1,2]. The often-discussed need for tools to assist decision makers has led to development of many clinical decision support (CDS) models and associated tools [3,4], capable of integrating disparate sources of information and guiding clinicians in their

decision-making task [5]. These tools have evolved from simple scoring systems to include complicated multivariate regression, neural networks, decision trees, and probabilistic models [6,7].

This paper focuses on a specific type of graphical probabilistic model, the *Bayesian network* (BN) [8]. BNs have become popular CDS models in medicine [9,10]. This popularity results from their ability to: (i) model complex problems with causal dependencies where a significant degree of uncertainty is involved; (ii) combine multiple sources of information including data and experts' judgement; (iii) present as an interpretable graphical structure; and, (iv) model interventions and reason both diagnostically and prognostically.

Early attempts to use Bayesian analysis for medical problems were considered unsuccessful because BN computation was intractable [11]. Development of efficient BN inference propagation algorithms [12,8]

\* Corresponding author.

E-mail address: [e.kyrimi@qmul.ac.uk](mailto:e.kyrimi@qmul.ac.uk) (E. Kyrimi).

<https://doi.org/10.1016/j.artmed.2021.102079>

Received 10 July 2020; Received in revised form 14 April 2021; Accepted 20 April 2021

Available online 22 April 2021

0933-3657/© 2021 Elsevier B.V. All rights reserved.

and continuing advances in computational power have made possible BNs capable of addressing real-world decision-support problems. This has motivated renewed research interest which has seen many thousands of publications proposing BN solutions for medical application [13]. However, despite the immense volume of published studies on medical BNs there is little evidence for their adoption in clinical practice as shown in our recent scoping review [14]. We found that the content of most literature concerning BNs developed as CDS focuses on BN development. Only a minority of published medical BN models directly contemplate the importance of and need for external validation, and fewer still pay any attention to BN usefulness and adoption in clinical practice. To the best of our knowledge, the chasm between research effort and clinical adoption has not drawn adequate attention [14,13]. This paper is one of a series of papers resulting from a large review conducted by our lab during 2019 investigating the development of BNs for use in healthcare. At first, a preliminary review of BNs in healthcare based only on a very small number of published papers was initially conducted by the authors [13] to identify potential research gaps and justify the conduct of this broader scoping review. Three papers resulted from our larger scoping review. While these papers are outputs using the same literature collection on BNs in healthcare, the contributions presented in each paper are entirely different such that there has been no repetition of findings across the three papers. In [15] the distribution of medical conditions observed in the studied literature has been explored. In [10] several unique objectives such as the aim of the published BNs, their development process, and their clinical usefulness have been studied.

Our earlier works first identified that while the literature presents many Bayesian-based clinical decision-support tools, none had seen adoption in front-line clinical care [13], and second that it was rare for the authors of published medical BNs to consider the process by which their solution might be implemented and used in clinical practice [10]. We contend that identifying the existence of these gaps created a need for additional investigation to identify the issues preventing adoption of BNs in healthcare, which is the primary aim of this paper. Motivated by the desire to enhance dialogue among researchers and gain deeper understanding for the neglected issue of BN adoption in practice, this paper focuses on exploring the benefits, barriers and facilitating factors for implementing BN-based solutions in clinical care as an approach to achieving that aim.

Acknowledging that a gap exists between developing an accurate model and demonstrating its clinical usefulness and impact on decision-making, several researchers have explored the stages necessary to implementing prediction models in clinical practice [16–18,7]. Additionally, several studies have investigated reasons why CDS models are not found in clinical practice [14,19]. We extend on these and review a collection of 116 recent papers proposing medical BNs for supporting clinical decision-making, with specific attention to identifying the benefits, barriers and facilitating factors. Benefits present as the positive outcomes that authors propose may result from implementation in clinical practice of their proposed BN-based system, such as improved patient outcomes or reduced costs and resource consumption. Benefits are usually realised by fixing one or more barriers through engagement of a facilitating factor. Barriers present as obstacles preventing clinical adoption of the BN, such as clinicians' reluctance or the high cost of implementation. Finally, facilitating factors are those elements that when applied, are able to overcome one or more existing barriers, and by doing so, promote successful implementation. Examples might include providing a user-friendly and contextual interface to minimise clinicians' reluctance to engage with new technologies.

In this paper we apply the ITPOSMO-BBF framework developed by McLachlan et al. [20] to explore the chasm between recently developed medical BNs and their lack of adoption in clinical practice. We present results of a literature review of works describing BNs in healthcare and identify the benefits, barriers and facilitating factors discussed by authors when proposing their BN-based CDS solution. The results of this

review can a) help identify reasons for the lack of adoption of BNs in healthcare; and b) promote efforts for clinical adoption of BN-based systems.

The remainder of this paper is organised as follows: Section 2 introduces background material regarding implementation of CDS model in clinical practice. Section 3 explains the ITPOSMO-BBF methodology used. Section 4 presents our results, while a discussion of our findings and a conclusion are provided in Section 5 and 6, respectively.

## 2. Background: implementation of clinical decision support systems

As a background to the work presented in this paper, this section describes the generally established framework for implementation of CDS systems and contextualises implementation of BN-based CDS within that framework. Based on the existing literature for implementing CDS systems [21,17,22,18] the following four main stages should be followed:

*Development:* A CDS model is developed by: (a) identifying important predictors; (b) verifying the assumptions behind the model; (c) estimating its predictive accuracy; (d) calibration; and (e) discrimination using the data used to develop the model, called *internal validation* [23]. It is a general requirement that key details that describe how a model was developed and validated be clearly reported to enable synthesis and critical appraisal of all relevant information [24]. Only with full and clear reporting of the model development process and internal validation can risk of bias and potential usefulness of models be adequately assessed [24–26].

*External validation:* After developing a CDS model it is crucial to validate its predictive accuracy on data not used during model development in a process called *external validation* [27,28]. External validation of model performance is crucial for verifying the model's generalisability. A clinical model is unlikely to be accepted if it has not been proven to work on disparate populations. External validation may use: (i) data collected from the same hospital but sampled from a later or earlier period, called *temporal validation*; data from different hospitals or countries, where the clinical care and definitions might be different, called *geographic validation*; or (iii) data sampled from demographically different individuals than those from which the model was developed, called *domain validation* [27,24]. If necessary, the model can be updated using knowledge gained from the validation process.

*Assessment of Clinical Usefulness:* A useful model should have an impact on clinical decision making. Assessment of the clinical usefulness of a model consists of three phases. *First*, the existing clinical situation and potential benefit of the tool must be clear [14]. Questions such as: (i) *What is the targeted population?* and (ii) *Who is going to use the model, at what point during the care and for what purpose?* should be clarified [29]. Also, we must investigate how the model will integrate into the clinical workflow and any barriers we might experience, such as ethical and regulatory restrictions [30,31]. *Second*, before conducting an extended impact analysis we must verify that the CDS model has potential to impact clinical decision-making [17,14,28,32]. The first objective at this phase is to verify the credibility, also known as face validity, of the model [21]. A model without credibility may not include well-known predictors for disease or treatment outcome and therefore clinicians might doubt its advice [14]. Thus, clinicians should review the model's logic, comprehension and relevance. After this, the potential impact of the CDS model should be examined [33] as this is the first sign for whether it is going to be used or not [22,18]. A system that rarely updates the user's decision will not be used, no matter how accurate it is. Another important element of usefulness considered during this phase is the model's usability. Clinicians must be able to use and comprehend the CDS system, as a complex system is unlikely to be used [34]. Hence, a user-friendly interface and easily interpretable predictions are vital [14]. It is also crucial to have a trustworthy and explainable model whose reasoning is clear to clinicians as they are unlikely to use a model

they do not understand or trust [35,36]. *The third and final phase* before implementation is undertaking an *actual impact* analysis. The impact analysis is an evaluation that can identify whether the CDS can sustain a long-term implementation, or if there are barriers that must still be resolved. The actual impact of a CDS system will differ from its potential impact. Clinicians will not always follow the system’s recommendations: they may not consult the system at all; apply it inaccurately; overrule its recommendations or be unable to implement its recommendations [17, 18].

*Adoption:* The final stage is the acceptance and implementation of the CDS to assist decision-making and guide patient management in clinical practice. Even though this stage represents the endpoint for development and implementation, the whole process should continue as new clinical knowledge is identified, thus operating like evidence-based medical practice, as a *learning cycle* [37]. New knowledge might include changes in the clinical guidelines used to define healthcare practice that should be evaluated for incorporation into the CDS.

### 3. Methodology

This work applies the ITPOSMO-BBF methodology [20] to a collection of literature proposing BNs for use in healthcare. We seek to provide a balanced comparison among benefits, barriers and facilitating factors as an approach to systematically assess the challenges that contribute to the lack of adoption for BNs in clinical practice. Our methodology involves three phases: Search and selection, Research framework, and Analysis.

#### 3.1. Search and selection

We used the literature collection selected from our recent scoping review on BNs in healthcare [10]. The search term used to derive the selected literature was:

“(((Bayes OR Bayesian) AND network) OR (probabilistic AND graphical AND model)) AND (medical OR clinical)”

Terms such as *Bayesian networks* or *graphical probabilistic models* were used here because they are widely observed in the targeted literature. Different ways for explaining the medical condition do occur, in that in some papers the exact condition is mentioned while in others broader terms such as *medical* or *clinical* accompanied by *application*, *condition*, or *setting*, are used. Our scoping review settled on the broader terms *medical* or *clinical* as they were commonly found in a larger collection of papers. Searching for specific medical conditions would have been impractical as there are many thousands of distinct known conditions.

As detailed in [10], due to the high number of articles returned,

further scrutiny was applied to narrow the collection to one which was reasonable to complete a defined review. This was achieved by assessing whether the described keywords were present in the title and abstract. Additional screening was conducted to exclude papers published outside the period 2012–2018, those that were not published in English, or those whose BN focus was not healthcare related. The remaining eligible papers were those possessing all the following identified characteristics in that they:

- 1 Describe a genuine BN model or BN adoption process
- 2 Are targeted clearly at a medical condition or application
- 3 Are intended to support clinicians or patients in decision making

#### 3.2. Research framework

Inspired by the research framework proposed by Yao et al. [38], the literature collection was reviewed using the logic process illustrated in Fig. 1. Benefits, barriers and facilitating factors for adopting BN-based systems in healthcare were identified using content analysis (CA) and thematic analysis (TA) [39]. In addition, the context in which authors described them, as well as their frequency in the collected literature was examined.

#### 3.3. Analysis

Authors often represented the same general idea in a variety of ways. CA and TA are separate but interrelated qualitative approaches for descriptive data analysis [39]. CA is an accepted systematic coding and categorisation method for investigating texts and resolving a quantitative description of the features [39,40]. CA establishes categories and then records the instances in which that category is evident or can be inferred from within the collected texts being analysed [40]. TA is a more qualitative method used to identify, analyse and report patterns, or themes, that emerge as being important within the material being analysed [39,40]. TA provides the systematic element characteristic of CA, while additionally affording the ability to combine analysis of frequency with analysis of *in context* meaning, therefore providing a more truly qualitative analysis [40]. CA and TA are established methods regularly used in clinical, nursing and other healthcare contexts [39,40]. While CA resolved many common concepts, it is only once TA was applied that these concepts could be described by their underlying contextual themes. The concepts and themes identified through these processes provided the necessary knowledge and data to support our ITPOSMO-BBF analysis.

One widely used approach for evaluating information technology implementation is ITPOSMO [41]. ITPOSMO stands for: (1) Information;




Fig. 1. Research framework.

(2) Technology; (3) Processes, (4) Objectives; (5) Staffing and skills; (6) Management systems and structures; and (7) Other resources, which includes such things as time and money. Heeks et al. [41] describe these *seven dimensions*, grouped into four *aspects*, (1) Information and Technology, (2) Process and Objectives and values, (3) Staffing and skills and Management, and (4) Other resources, as being capable of exploring the gap between a system’s design and the reality of its implementation. The ITPOSMO framework was initially proposed for evaluation of e-government projects [41,42], but has also been used to investigate how technology can improve healthcare practices [38] or identify problems in cloud health information system projects [43]. Recently, an extension, ITPOSMO-BBF, was presented for comparative analysis of barriers, benefits, and facilitators in health information technology: and was specifically applied to evaluation of *electronic health records* (EHR) and *learning health systems* (LHS) implementations [20].

Advantages for using the ITPOSMO-BBF framework in this study are that; (1) apart from simply identifying barriers and facilitating factors for adopting BN-based systems in healthcare, we can also quantify their relationships; (2) we can identify benefits when adopting BN-based systems in healthcare; and (3) while ITPOSMO was developed as a retrospective analysis of completed projects, ITPOSMO-BBF can be used to understand, plan for, and mitigate potential barriers prior to adoption. The last advantage is especially important in this study as most published BN models are developed with the potential to be used in clinical practice, but adoption in clinical settings remains elusive.

For each of the four ITPOSMO-BBF aspects, the frequency of (1) benefits; (2) facilitating factors; and (3) barriers in the studied literature are presented following the layout illustrated in Fig. 2. The (4) relationship between a barrier and a facilitator is also provided using a weighted line, with the thickness of the line indicating the number of authors who identified that particular relationship.

#### 4. Results

##### 4.1. Results of literature search

The literature collection was drawn from our recent scoping review on BNs in healthcare [10]. As shown in Fig. 3, the search initially identified 3810 papers. In addition to the exclusions described in Section 3.1, we excluded papers that were focused on Bayesian statistics or meta-analyses rather than BNs. Graphical models, such as naive BNs, which are the simplest BNs and structurally assume all variables are “conditionally independent” (meaning that they are independent once the classification is known), and other similar approaches including neural networks, were also excluded. This resulted in 123 papers for inclusion [10]. However, after an initial reading of the remaining papers a further 7 were excluded that failed to provide sufficient information for assessing benefits, barriers or facilitating factors for adopting their

BN model in clinical practice. Four of these were purely methodological, two were survey papers, and the final paper had been retracted by the publishing journal who noted acknowledgement of the authors that it contained errors. Thus, the final literature collection for this work contains 116 papers describing 111 unique BN-based CDS systems, and it can be found in [https://pambayesian.org/wp-content/uploads/2020/06/BBF\\_Results.xlsx](https://pambayesian.org/wp-content/uploads/2020/06/BBF_Results.xlsx).

There was a large distribution of medical conditions discussed in the studied literature. Cardiac conditions, cancer, psychological and psychiatric disorders, and lung and breathing disorders made up 59 % of the medical conditions. The remaining 41 % of papers were spread across a diverse collection of seemingly random topics. A more detailed description of the medical conditions discussed in the literature can be found at [15]. The main type of reasoning identified in the studied literature was observational reasoning. More specifically, in 90 of the papers reasoning from observations was performed. Diagnosis of a medical condition given a set of known risk factors and observed signs and symptoms was the most common decision type among those papers. Finally, in 37 papers, the BN aim was to learn associations or causal relationships from data in order to understand and provide causal attribution to a disease. For a review and detailed description of the decision types and reasoning strategies identified from our literature collection, we refer the reader to [10].

##### 4.2. Content and thematic analysis

CA and TA were used to identify benefits, barriers and facilitating factors that impact or support adoption of BN-based systems in healthcare. Using TA four main themes were identified within the collected literature; (1) data, (2) resource, (3) resistance, and (4) performance. Each theme clusters a set of related contents. The contents corresponding to each theme are described in Table 1 and the Appendix. As observed in Table 1 contents are divided regarding whether they have been encountered as benefits, barriers or facilitating factors. For instance, on the one hand, Velikova et al. [44] mention that using a BN-based CDS as a home-monitoring system has the benefit of enhancing data collection as it gives access to a large collection of clinical and laboratory data. On the other hand, Vemulapalli et al. [45] mention the same content of enhanced data collection as a facilitating factor for overcoming the barrier of not having sufficient or well-structured patient-centric healthcare data.

##### 4.3. ITPOSMO-BBF framework

This section presents results of our analysis of the literature, including the frequency with which authors discussed themes in context, as described in Table 1. The themes and contexts are mapped to the four ITPOSMO aspects (1) Information and Technology (IT), (2) Process and




Fig. 2. Visualisation of the ITPOSMO analysis.




Fig. 3. Literature selection.

Objectives and values (PO), (3) Staffing and skills and Management (SM), and (4) Other resources (O).

#### 4.3.1. Information and technology

A number of IT barriers were described and are shown in Fig. 4. Several authors reported that quality and availability of medical data was lacking [46–49]. Regardless of the efforts on overcoming lack of data when developing BNs using medical expertise and/or literature (an important advantage of BNs over purely data-driven models) the poor quality and/or lack of medical data remain an issue for the model's performance and usability [50–52]. Incomplete data can result from numerous causes not limited to cost limitations with regards to equipment or inadequate assessment of patient-related factors [53,54,49]. However, even if data is available it may exist unstructured [55–57]. Different countries, or even hospitals within countries, store data using different classification standards which complicates integration and aggregation of patient datasets [46,58,48]. Another barrier of BN-based systems was their limited capability to address continuous data. Data

discretisation is often required, which does not generally follow clinical reasoning and can result in information loss [59–62]. Finally, BN-based systems usually rely on non-randomised observational data. True causal relationships such as those observed between drugs and adverse events are not easily detected from observational data [63–66].

Several authors identified that government incentives to transition from paper-based patient information to electronic medical records improves availability of healthcare data while also providing an opportunity to implement BN-based systems [50,45,67]. Additionally, standardised note templates with predefined fields and responses can enable relevant structured data to be populated directly into clinical databases [55,57,48]. Richer datasets can support higher predictive capability, delivering superior and more useful CDS systems [68,46,69,70]. Data availability can also be improved by developing technologies for healthcare providers to share information between facilities [48,67]. In addition, integrating different sources of information with experts judgement can sometimes overcome limitations in the data alone [59,56,68]. Advances in software tools to better deal with continuous data

**Table 1**  
Themes and contexts identified in the selected literature.

Themes	Contexts		
	Benefits	Barriers	Facilitating Factors
Data	enhance data collection,deal with problematic data,use various source of information, improve data exchange	Availability/ quality,lack of randomisation, unstructured,not standardised, discretisation	enhance data collection, information sharing among clinicians,use various sources of information
Resource	cost saving		improvements in algorithms and software, access to technology,user-friendly interface
Resistance	easy to understand,applicable in medical situations with uncertainty, predictions in the presence of missing data,incremental data entry, confidence in the prediction,flexibility,query any given node	clinicians' resistance,lack of credibility, lack of clinical impact	Explanation of reasoning,clinical involvement,follow current practice, impact analysis
Performance	improved accuracy,improve quality of care,care process standardisation, risk understanding,multiple reasoning approaches, accelerate clinical decisions,assist clinical decisions,improve patient outcome,fit constantly evolving information, explore numerous causal relationships	poor predictive accuracy,lack of generalisability	internal validation,validate in different populations

and enable access to faster and more flexible technology can overcome barriers of data discretisation and make BN-based systems more accessible [71,62,72,67,44,73].

Application of these facilitating factors to resolve the barriers identified earlier may give rise to numerous benefits described by many authors. Adoption of a BN-based systems in practice may enhance data collection. For example, BN-based systems can be used to systematically collect important clinical data [59,55,44]. BN-based systems can also be used to make health information more widely accessible [74,55]. BN-based systems are able to deal with noisy, incomplete and heterogeneous data [75,76,57,49,77] and perform reasoning using data from multiple sources, including historical data, sensors, and experts [78,44,47,73]. This capability to handle heterogeneous information is essential to delivering precision medicine [68].

#### 4.3.2. Process and objectives

Lack of clinical impact, shown in Fig. 5, was the main PO barrier for acceptance of BN-based systems in healthcare [79]. Evaluating the system's impact on clinical decision making, its usefulness in practice, and the degree of practitioner acceptance were considered important and necessary steps to aid adoption in practice [80,69,62,81]. Further, BN-based systems should adhere with and be integrated into current clinical practice in order to be accepted by clinicians [72,82,47].

Even if PO barriers and facilitating factors are largely ignored, many benefits were discussed in the literature. BN-based systems are suitable tools in helping medical-related problems such as diagnosis, prognosis and treatment selection due to the uncertain nature of these problems and the BN's ability to efficiently deal with this uncertainty [83,79,84–86,49,81]. The BNs ability to perform more than just reasoning from evidence makes it a valuable tool in clinical decision making [71,47,50]. Another important benefit is that they can update their prediction as new patient information becomes available [72,50,87,85]. As clinical knowledge and data increase and new standards for evidence-based clinical practice are implemented, BN-based systems can be updated to maintain relevance [50,88,84,89]. This continual apprenticeship makes it possible to refine the predictive quality of the model and adapt medical practices [74]. Moreover, BN-based systems can enhance clinical objectives, such as aiding and accelerating clinical decision making [71,82,70,81,90,69,91], improving understanding of significant associated risk [92,93], improving patient outcomes [71,48,65] and improving overall quality of care [90,73]. In this way BN-based systems can standardise medical care by providing a consistent standard of advice [74].

#### 4.3.3. Skills and management

Clinicians' resistance to health IT systems was the most impactful of the two SM barriers discussed in the literature and shown in Fig. 6. Attaining cooperation and acceptance of clinicians is crucial for wider

adoption of any health IT, but especially BN-based systems. Although progress has been made, BN-based tools remain rarely used in clinical practice, possibly because the medical field tends to be cautious about adopting new technology unless its advantages are entirely clear [44, 88]. BNs are not the first modelling technique considered by health policymakers who are generally more familiar with regression models [65]. Another barrier to widespread adoption of BN-based systems is the requirement for extensive development effort and subjectivity involved when relying on clinicians' judgement [69]. Clinicians can also resist processes that may interfere with their daily workflow or challenge their autonomy [47]. And finally, another barrier related to clinician's resistance arises when a model lacks credibility, also known as face validity [44,53,54,79,80].

Most effort aimed at facilitating adoption of BN-based systems has gone towards resolving clinicians' resistance. Clinical involvement was considered an important facilitating factor for resolving both clinicians' resistance and the lack of credibility. Cooperating with multiple clinicians has been considered beneficial to developing a credible BN-based system that captures medical knowledge, property desired by clinicians [83,44,94,54,69]. BN-based systems should be reviewed by clinicians in order to obtain their input and agreement if we are to achieve adoption in practice [80,65,95]. In addition, development of a user-friendly interface was described as important element for assisting adoption [59,71,96,87,97]. A BN-based system's interpretability was also mentioned as a facilitator for overcoming clinicians' resistance. Their graphical structure can make them more comprehensible, which is important in medical applications [67,54,58]. An explanation of the system's prediction could also be beneficial [44,92]. The fact that BN-based systems can be easily understood is also one of their main benefits [98,65,97,99].

#### 4.3.4. Other resources

The other resources and constraints (O) barriers and facilitating factors, shown in Fig. 7, contain those attributes that did not naturally fall within the other ITPOSMO aspects, and which are mainly about the system's overall performance. A model with poor predictive accuracy cannot be applied in clinical practice [79,64,96]. Validating the system's performance was considered as a necessary facilitating factor for assisting its adoption [71,46,100,86]. In addition, even if the BN-based system has a good predictive performance, it will fail to be accepted in practice, if it has no generalisability [79,101,65]. Thus, external validation must be undertaken in different cohorts and geographical areas before the BN-based system can be used routinely in clinical practice [102,48,73,103,81].

Several benefits were mentioned in the literature. Using BN-based systems to optimise processes at all stages of care, as explained in PO benefits, can significantly benefit with respect to healthcare cost [63,45,49,104]. BN-based systems, as opposed to deterministic approaches,




Fig. 4. Barriers, facilitating factors and benefits regarding Information and Technology (ITPOSMO).




Fig. 5. Barriers, facilitating factors and benefits regarding Process and Objectives (ITPOSMO).

also provide confidence in the prediction [78,79,105,55], resulting in a greater flexibility [101]. It is also possible to query any given node in the BN, making BN-based systems substantially more useful in clinical practice when compared to models constructed with reliance on specific outcome variables [47,82,106]. Further, unlike logistic regressions and

other modelling techniques, BN-based systems are not limited to linear relationships. In contrast, they are able to model complex relationships between variables when conditions of causality and conditional independence are involved, which is of significant benefit in clinical decision making [101,48,107,106]. Finally, BN-based systems have a reputation




Fig. 6. Barriers, facilitating factors and benefits regarding Skills and Management (ITPOSMO).




Fig. 7. Barriers, facilitating factors and benefits regarding Other resources (ITPOSMO).



for achieving accurate results [79,58,102,108,100].

## 5. Discussion

Despite the large volume of published studies on medical BNs, there is little evidence for their adoption in clinical practice. There has been little attention paid to the gap between developing an accurate BN and demonstrating its clinical usefulness for decision making. In this paper we have provided a deeper insight into the gap. Using the ITPOSMSO-BBF framework we have identified the *benefits*, *barriers* and *facilitating factors* discussed in the literature for adopting BN-based systems in clinical practice. This is an important study because most published BN models are developed with potential for positive effect if applied in clinical practice, yet their adoption remains elusive.

The barriers discussed in the literature that seem to obstruct BN-based system adoption in clinical practice can be grouped into: (1) data inadequacies; (2) clinicians' resistance to new technologies; (3) BN-based systems that lack of clinical credibility; (4) failure to demonstrate clinical impact; (5) absence of an acceptable predictive performance; and (6) absence of evidence for a model's generalisability. Similarly, the facilitating factors mentioned in the studied literature for overcoming existing barriers and enabling adoption can be categorised into: (1) data collection improvements; (2) software and technological improvements; (3) having interpretable and easy to use BN-based systems; (4) clinical involvement in the development or review of the model; (5) investigation of model's clinical impact; (6) internal validation of the model's performance; and (7) external validation of the model.

As presented in Section 4, the frequency with which authors discussed the identified barriers and facilitating factors was quite low. Even rarer were authors who actually connected some form of facilitating factor with the barriers they had identified. For instance, having a user-friendly interface was described as a necessary step for adopting a BN-based system in practice, yet this was rarely reported as a facilitating factor for reducing the barrier of clinicians' resistance. This lack of more profound thinking that might link the barriers faced when implementing BN-based systems in practice with solutions to overcome those barriers is very likely the reason we see such a chasm between research effort and adoption in practice.

While the findings presented in this paper result from applying the established ITPOSMSO-BBF framework as part of a well-structured review of a large number of papers, it does not come without its limitations:

- 1 *Possible missing relevant papers*: Even if the studied literature is a representative sample of papers published in both medical and AI journals and conference proceedings, it may not reflect the entire range of the literature regarding BNs in healthcare. We looked for keywords such as *Bayesian Networks*, *probabilistic graphical models*, *medical*, *clinical* to appear in the abstract of each paper. It is possible that a number of relevant papers were not included because they did not use the selected keywords in their abstract. This is especially true in cases where the actual name of the medical condition is described without mentioning the words *medical* or *clinical*. However, we believe that the large number of selected papers was sufficient for drawing conclusions.
- 2 *Limited time period*: Because of the plethora of published papers and the time needed for review and data extraction we limited the study to the seven-year period 2012–2018. Although this is a safe number of years to be able to draw conclusions, papers published before 2012 were not reviewed. As the literature search was conducted in early 2019 as part of a larger review, papers published in 2019 and 2020 are not included in the collection. More recent publications might

focus more on BN adoption, describing better the associated benefits, barriers and facilitating factors, as this is a topic rising more and more attention in the research community. However, we do not believe that omitting papers published in 2019–2020 impacts significantly on the findings of this review.

- 3 *Possible missing evidence of adoption*: It is possible that evidence for subsequent translation and adoption of BNs into the real world may well appear in places other than academic journals, such as company reports, media, and general commentaries. For instance, based on limited marketing materials, there are indications that Babylon is using BNs in their non-public research and application development [109], including what is claimed as the largest BN in the world [110], while clients of BN tools companies like Hugin may be using BN medical decision support systems like Munin [111,112]. As a result, BNs that have been used in practice as well as the encountered benefits, barriers and facilitating factors, but have not been published in scientific journals might have been missed. It is, of course, also possible that real use of medical BNs is deliberately not published for reasons of either commercial or patient confidentiality.

## 6. Conclusion

The barriers that can prevent BNs from being adopted in clinical practice can be broadly grouped into: lack of resources, clinical resistance, insufficient clinical impact and model performance. While very few authors demonstrate understanding of these barriers to BN adoption, the many benefits that could arise from implementing BNs were thoroughly discussed in the literature. Most discussed benefits fell within the PO aspect and concerned the positive contribution BNs could make to meeting clinical objectives, such as assisting clinical decisions, improving understanding of associated risks, or improving the overall quality of care. The PO aspect is a prime example of the most significant issue in BN literature: that while many benefits can be identified by authors, the barriers limiting our ability to realise those benefits, and any facilitating factors to aid mitigation of barriers, are subjects that both remain notably neglected.

There is a clear gap between the development of such models and the ability to implement a useful decision-support tool into clinical practice. Reasons for this chasm include an overemphasis on technical aspects of algorithm development, with insufficient attention to improving the usability, explainability, trust and potential impact of these models. More attention needs to be given to the process of translating accurate prediction models into useful, usable and explainable CDS systems if the potential benefits of these are to be realised. Thus, we recommend BN developers should move from “why does it fail” towards “how to succeed” and address the barriers earlier to avoid the adoption challenges continuing [113]. The key to this is getting the *right* information, to the *right* people, in the *right* format, through the *right* channels, at the *right* times to enhance decisions [114].

## Acknowledgements

EK, SM, MRN, AF, WM and NF acknowledge support from the Engineering and Physical Sciences Research Council (EPSRC) under project EP/P009964/1: PAMBAYESIAN: Patient Managed decision-support using Bayes Networks. EK and WM acknowledge also support from the US Defense Medical Research and Development program under COMBAT-AID project: COMPUTER Battlefield Assistance in Trauma care and injury Decision-support (W81XWH1920047). KD acknowledges support from the School of Fundamental Sciences, Massey University, New Zealand, for his study sabbatical and visits with the PamBayesian team.

## Appendix A

An extension of Table 1 in which each theme and context are connected to specific literature identified using the row number in [https://pambayesian.org/wp-content/uploads/2020/06/BBF\\_Results.xlsx](https://pambayesian.org/wp-content/uploads/2020/06/BBF_Results.xlsx).

Themes	Contexts					
	Benefits	Barriers	Facilitating Factors			
Data	Enhance data collection, deal with problematic data, use various source of information, improve data exchange	Rows: 4, 10, 16, 20, 26, 31, 37, 48, 52, 55, 57, 61, 74, 76, 82, 87, 91, 97, 100, 114, 117	Availability/ quality, lack of randomisation, unstructured, not standardised, discretisation	Rows: 4–5, 7, 9–12, 15–17, 24–25, 30, 34, 38, 4, 49, 53, 56–57, 59–60, 63–64, 68, 74–75, 82, 87, 90–91, 93, 95, 97, 104–105, 108–110, 114, 119	enhance data collection, information sharing among clinicians, use various source of information	Rows: 4, 5, 10, 16, 17, 44, 60, 65, 85, 91, 93, 101, 110, 114
Resource	cost saving	Rows: 9, 11, 23, 24, 68, 100, 101			improvements in algorithms and software, access to technology, user-friendly interface	Rows: 4, 6, 8, 10, 14, 17, 30, 64, 70, 74, 78, 85, 97, 98, 100, 102, 107, 113
Resistance	Easy to understand, applicable in medical situations with uncertainty, provide predictions in the presence of missing data, incremental data entry, confidence in the prediction, flexibility, query any given node	Rows: 4–6, 9–16, 18, 20–21, 23 25–27, 29, 32, 36–37, 39, 41, 43–46, 48–57, 59–60, 62–64, 66, 70–72, 74–78, 80, 82–87, 89, 91, 93–100, 102–103, 107, 110–114	clinicians' resistance, lack of credibility, lack of clinical impact	Rows: 9, 11, 15, 16, 30, 41, 60, 80, 87, 100, 108	Explanation of reasoning, clinical involvement, follow current practice, impact analysis	Rows: 8, 9, 13, 15, 16, 27, 30, 45, 59, 64, 70, 78, 81, 84, 85, 87, 98, 99, 100, 102, 107
Performance	improved accuracy, improve quality of care, care process standardisation, risk understanding, multiple reasoning approaches, accelerate clinical decisions, assist clinical decisions, improve patient outcome, fit constantly evolving information, explore causal relationships	Rows: 4–16, 20, 23–27, 29–46, 48–51, 53–60, 62–64, 66–87, 91–105, 107, 110, 112–115	poor predictive accuracy, lack of generalisability	Rows: 14, 15, 41, 49, 103, 115	internal validation, validate in different populations	Rows: 5, 6, 7, 10, 16, 20, 25, 26, 27, 33, 40, 44, 46, 63, 71, 75, 78, 84, 95, 97, 104, 114

## References

- [1] Tversky A, Kahneman D. Judgment under uncertainty: heuristics and biases. *Sci. New Ser.* 1974;185(4157):1124–31. <https://doi.org/10.1126/science.185.4157.1124>.
- [2] Bornstein BH, Emler AC. Rationality in medical decision making: A review of the literature on doctors' decision-making biases. *J Eval Clin Pract* 2001;7(2): 97–107. <https://doi.org/10.1046/j.1365-2753.2001.00284.x>.
- [3] Patel VL, et al. The coming of age of artificial intelligence in medicine. *Artif Intell Med* 2009;46(1):5–17. <https://doi.org/10.1016/j.artmed.2008.07.017>.
- [4] Shortliffe EH, Cimino JJ. *Biomedical informatics: Computer applications in health care and biomedicine*. 4th ed. 2013.
- [5] Druzdzel MJ, Flynn RR. *Decision support systems*. 2nd ed. *Encyclopedia of Library and Information Science*; 2002.
- [6] Lucas PJF, Abu-Hanna A. Prognostic methods in medicine. *Artif Intell Med* 1999; 15:105–19.
- [7] Adams ST, Leveson SH. Clinical prediction rules. *BMJ* 2012;344:d8312. <https://doi.org/10.1136/bmj.d8312>.
- [8] Pearl J. *Probabilistic reasoning in intelligent systems: networks of plausible inference*. Morgan Kaufmann Publishers Inc.; 1988.
- [9] Lucas P, van der Gaag LC, Abu-Hanna A. Bayesian networks in biomedicine and health-care. *Artif Intell Med* 2004;30(March):201–14. <https://doi.org/10.1016/j.artmed.2003.11.001>.
- [10] Kyrimi E, McLachlan S, Dube K, Neves MR, Fahmi A, Fenton N. A comprehensive scoping review of Bayesian networks in healthcare: past, present and future. *ArXiv* 2020. doi: 2002.08627.
- [11] Cooper GF. The computational complexity of probabilistic inference using Bayesian belief networks. *Artif Intell* 1990;42(2–3):393–405. [https://doi.org/10.1016/0004-3702\(90\)90060-D](https://doi.org/10.1016/0004-3702(90)90060-D).
- [12] Lauritzen SL, Spiegelhalter DJ. Local computations with probabilities on graphical structures and their application to expert systems. *J R Stat Soc Ser B* 1988;50:157–224. <https://doi.org/10.1007/BF01531016>.
- [13] Kyrimi E, McLachlan S, Dube K, Fenton N. Bayesian networks in healthcare: the chasm between research enthusiasm and clinical adoption. *medRxiv Prepr.* 2020. <https://doi.org/10.1101/2020.06.04.20122911>.
- [14] Kleinrouweler CE, et al. Prognostic models in obstetrics: available, but far from applicable. *Am J Obstet Gynecol* 2015;214(1):79–90. <https://doi.org/10.1016/j.ajog.2015.06.013>. e36.
- [15] McLachlan S, Dube K, Hitman GA, Fenton NE, Kyrimi E. Bayesian networks in healthcare: distribution by medical condition. *Artif Intell Med* 2020;107 (January):101912. <https://doi.org/10.1016/j.artmed.2020.101912>.
- [16] Wyatt JC, Altman DG. Commentary: Prognostic models: clinically useful or quickly forgotten? *BMJ* 1995;311(December):1539–41. <https://doi.org/10.1136/bmj.311.7019.1539>.
- [17] Reilly BM, Evans AT. Translating clinical research into clinical practice: impact of using prediction rules to make decisions. *Ann Intern Med* 2006;144:201–9 [Online]. Available: <http://www.scopus.com/inward/record.uri?eid=2-s2.0-33644855006&partnerID=40&md5=903fb2ad9598fcbdc1984db9a9af8784>.
- [18] Moons KGM, Altman DG, Vergouwe Y, Royston P. Prognosis and prognostic research: application and impact of prognostic models in clinical practice. *BMJ* 2009;338(January):1487–90. <https://doi.org/10.1136/bmj.b606>.
- [19] Sendak M, et al. The human body is a black box': supporting clinical decision-making with deep learning. *Proceedings of ACM Conference on Fairness, Accountability, and Transparency (FAT\* 2020)* 2020;99–109. <https://doi.org/10.1145/3351095.3372827>.
- [20] McLachlan S, et al. A framework for analysing learning health systems: are we removing the most impactful barriers? *Learn. Heal. Syst.* 2019;3(4):1–15. <https://doi.org/10.1002/lrh2.10189>.
- [21] Blackmore CC. Clinical prediction rules in trauma imaging: Who, how, and why? *Radiology* 2005;235:371–4. <https://doi.org/10.1148/radiol.2352040307>.
- [22] Toll DB, Janssen KJM, Vergouwe Y, Moons KGM. Validation, updating and impact of clinical prediction rules: a review. *J Clin Epidemiol* 2008;61(November): 1085–94. <https://doi.org/10.1016/j.jclinepi.2008.04.008>.
- [23] Steyerberg EW, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010;21(1 January):128–38. <https://doi.org/10.1097/EDE.0b013e3181c30fb2>.
- [24] Collins GS, Reitsma JB, Altman DG, Moons KGM. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): The TRIPOD statement. *Ann Intern Med* 2015;162:55–63. <https://doi.org/10.7326/M14-0697>.

- [25] Bouwmeester W, et al. Reporting and methods in clinical prediction research: a systematic review. *PLoS Med* 2012;9(5 January):1–12. <https://doi.org/10.1371/journal.pmed.1001221>.
- [26] Heus P, et al. Uniformity in measuring adherence to reporting guidelines: the example of TRIPOD for assessing completeness of reporting of prediction model studies. *BMJ Open* 2019;9(4):1–6. <https://doi.org/10.1136/bmjopen-2018-025611>.
- [27] Moons KGM, et al. Risk prediction models: II. External validation, model updating, and impact assessment. *Heart* 2012;98(March):691–8. <https://doi.org/10.1136/heartjnl-2011-301247>.
- [28] Park SH, Han K. Methodologic guide for evaluating clinical performance and effect of artificial intelligence technology for medical diagnosis and prediction. *Radiology* 2018;286(3):800–9. <https://doi.org/10.1148/radiol.2017171920>.
- [29] Moons KGM, et al. PROBAST: a tool to assess risk of bias and applicability of prediction model studies: explanation and elaboration. *Ann Intern Med* 2019;170(1):W1–33. <https://doi.org/10.7326/M18-1377>.
- [30] Wallace E, et al. Framework for the impact analysis and implementation of Clinical Prediction Rules (CPRs). *BMC Med Inform Decis Mak* 2011;11(1 January): 62. <https://doi.org/10.1186/1472-6947-11-62>.
- [31] Reddy S, Allan S, Coghlan S, Cooper P. A governance model for the application of AI in health care. *J Am Med Inform Assoc* 2020;27(3):491–7. <https://doi.org/10.1093/jamia/ocz192>.
- [32] Neves MR, Marsh DWR. Modelling the impact of AI for clinical decision support. *Conference on Artificial Intelligence in Medicine in Europe* 2019:292–7.
- [33] Reilly BM, et al. Impact of a clinical decision rule in the emergency department. *Emerg Med (N Y)* 2002;288(3).
- [34] Grady D, Berkowitz Sa. Why is a good clinical prediction rule so hard to find? *Arch Intern Med* 2011;171(19):1701–2. <https://doi.org/10.1001/archinternmed.2011.482>.
- [35] Lacave C, Díez FJ. A review of explanation methods for Bayesian networks. *Knowl Eng Rev* 2002;17(2 April):107–27.
- [36] Kyrimi E, Mossadegh S, Tai N, Marsh W. An incremental explanation of inference in Bayesian networks for increasing model trustworthiness and supporting clinical decision making. *Artif Intell Med* 2020;103(August):101812. <https://doi.org/10.1016/j.artmed.2020.101812>. 2019.
- [37] Buysse V, Winton P, Rous B, Epstein D, Lim C-I. Evidence-based practice: foundation for the CONNECT 5-step learning cycle in professional development. *Zero Three* 2012;32(4):25–9.
- [38] Yao W, Chu CH, Li Z. The use of RFID in healthcare: benefits and barriers. 2010 IEEE International Conference on RFID-Technology and Applications, RFID-TA 2010, 2010 10:128–34. <https://doi.org/10.1109/RFID-TA.2010.5529874>. no. July.
- [39] Vaismoradi M, Turunen H, Bondas T. Content analysis and thematic analysis: implications for conducting a qualitative descriptive study. *Nurs. Heal. Sci.* 2013; 15(3):398–405. <https://doi.org/10.1111/nhs.12048>.
- [40] Joffe H, Yardley L. Content and thematic analysis. *Res. Methods Clin. Heal. Psychol.* 2004;56.
- [41] Heeks R, Mundy D, Salazar A. Why health care information systems succeed or fail. *Inf. Syst. Public Sect. Manag. Work. Pap. Ser.* 1999;(9):1–25 [Online]. Available: [http://www.man.ac.uk/idpm/idpm\\_dp.htm#isps\\_wp](http://www.man.ac.uk/idpm/idpm_dp.htm#isps_wp).
- [42] Rughatjaroen K. Success of electronic government project in Bangkok metropolis: an ITPOSMO approach. *Int J Soc Sci Humanit Invent* 2015;5(9): 783–7. <https://doi.org/10.7763/ijssh.2015.v5.556>.
- [43] Syamsuddin I. Novel gap analysis framework for cloud health information systems. *J Theor Appl Inf Technol* 2016;87(3):415–21.
- [44] Velikova M, Van Scheltinga JT, Lucas PJF, Spaanderman M. Exploiting causal functional relationships in Bayesian network modelling for personalised healthcare. *Int J Approx Reason* 2014;55:59–73. <https://doi.org/10.1016/j.ijar.2013.03.016>.
- [45] Vemulapalli V, et al. Non-obvious correlations to disease management unraveled by Bayesian artificial intelligence analyses of CMS data. *Artif Intell Med* 2016;74: 1–8. <https://doi.org/10.1016/j.artmed.2016.11.001>.
- [46] Akhtar MJ, Utne IB. Human fatigue's effect on the risk of maritime groundings - A Bayesian Network modeling approach. *Saf Sci* 2014;62:427–40. <https://doi.org/10.1016/j.ssci.2013.10.002>.
- [47] Sesen MB, et al. Lung cancer assistant: a hybrid clinical decision support application for lung cancer care. *J R Soc Interface* 2014;11:20140534. <https://doi.org/10.1098/rsif.2014.0534>.
- [48] Jochems A, et al. Distributed learning: developing a predictive model based on data from multiple hospitals without data leaving the hospital – a real life proof of concept. *Radiother Oncol* 2016;121(3):459–67. <https://doi.org/10.1016/j.radonc.2016.10.002>.
- [49] Bukhanov N, Balakhontceva M, Kovalchuk S, Zvartau N, Konradi A. Multiscale modeling of comorbidity relations in hypertensive outpatients. *Procedia Comput Sci* 2017;121:446–50. <https://doi.org/10.1016/j.procs.2017.11.060>.
- [50] Loghmanpour NA, Druzzdel MJ, Antaki JF. Cardiac health risk stratification system (CHRiSS): a Bayesian-based decision support system for left ventricular assist device (LVAD) therapy. *PLoS One* 2014;9(11):1–10. <https://doi.org/10.1371/journal.pone.0111264>.
- [51] Ojeme B, Mbogho A. Selecting learning algorithms for simultaneous identification of depression and comorbid disorders. *Procedia Comput Sci* 2016;96:1294–303. <https://doi.org/10.1016/j.procs.2016.08.174>.
- [52] HeidarAbadi NN, Hakemi L, Kolivand P, Safdari R, Saeidi MG. Comparing performances of intelligent classifier algorithms for predicting type of pain in patients with spinal cord injury. *Electron Physician* 2017;9(9):4847–52. <https://doi.org/10.19082/5212>.
- [53] Yet B, Perkins Z, Fenton N, Tai N, Marsh W. Not just data: A method for improving prediction with knowledge. *J Biomed Inform* 2014;48(April):28–37. <https://doi.org/10.1016/j.jbi.2013.10.012> [doi].
- [54] Bandyopadhyay S, et al. Data mining for censored time-to-event data: a Bayesian network model for predicting cardiovascular risk from electronic health record data. *Data Min Knowl Discov* 2015;29(4):1033–69. <https://doi.org/10.1007/s10618-014-0386-6>.
- [55] Agarwal P, Verma R, Mallik A. Ontology based disease diagnosis system with probabilistic inference. 2016. <https://doi.org/10.1109/IICIP.2016.7975383>.
- [56] Constantinou AC, Fenton N, Marsh W, Radlinski L. From complex questionnaire and interviewing data to intelligent Bayesian network models for medical decision support. *Artif Intell Med* 2016;67:75–93. <https://doi.org/10.1016/j.artmed.2016.01.002>.
- [57] Solomon NP, Dietsch AM, Dietrich-Burns KE, Styrmisdottir EL, Armao CS. Dysphagia management and research in an acute-care military treatment facility: the role of applied informatics. *Mil Med* 2016;181(5S):138–44. <https://doi.org/10.7205/milmed-d-15-00170>.
- [58] Caillet P, Klemm S, Ducher M, Aussem A, Schott AM. Hip fracture in the elderly: a re-analysis of the EPIDOS study with causal Bayesian networks. *PLoS One* 2015; 10(3):1–12. <https://doi.org/10.1371/journal.pone.0120125>.
- [59] Berchiolla P, et al. Predicting severity of pathological scarring due to burn injuries: a clinical decision making tool using Bayesian networks. *Int Wound J* 2012;11(3):246–52. <https://doi.org/10.1111/j.1742-481X.2012.01080.x>.
- [60] Merli M, Moscatelli M, Mariotti G, Pagliaro U, Bernardelli F, Nieri M. A minimally invasive technique for lateral maxillary sinus floor elevation: a Bayesian network study. *Clin Oral Implants Res* 2016;27(3):273–81. <https://doi.org/10.1111/cir.12524>.
- [61] Luo Y, et al. Unraveling biophysical interactions of radiation pneumonitis in non-small-cell lung cancer via Bayesian network analysis. *Radiother Oncol* 2017;123(1):85–92. <https://doi.org/10.1016/j.radonc.2017.02.004>.
- [62] Magrini A, Luciani D, Stefanini FM. A probabilistic network for the diagnosis of acute cardiopulmonary diseases. *Biom J* 2018;60(1):174–95. <https://doi.org/10.1002/bimj.201600206>.
- [63] Duan L, Khoshsheshin M, Street WN, Liu M. Adverse drug effect detection. *IEEE J Biomed. Heal. Informatics* 2013;17(2):305–11. <https://doi.org/10.1109/TTB.2012.2227272>.
- [64] Kim M, Cheeti A, Yoo C, Choo M, Paick JS, Oh SJ. Non-invasive clinical parameters for the prediction of urodynamic bladder outlet obstruction: analysis using causal Bayesian networks. *PLoS One* 2014;9(11):9–14. <https://doi.org/10.1371/journal.pone.0113131>.
- [65] Chao YS, et al. A network perspective on patient experiences and health status: the medical expenditure panel survey 2004 to 2011. *BMC Health Serv Res* 2017; 17(1):1–12. <https://doi.org/10.1186/s12913-017-2496-5>.
- [66] Xu S, et al. Modeling interrelationships between health behaviors in overweight breast cancer survivors: applying Bayesian networks. *PLoS One* 2018;13(9):1–13. <https://doi.org/10.1371/journal.pone.0202923>.
- [67] Seixas FL, Zadrozny B, Laks J, Conci A, Muchaluat Saade DC. A Bayesian network decision model for supporting the diagnosis of dementia, Alzheimer's disease and mild cognitive impairment. *Comput Biol Med* 2014;51:140–58. <https://doi.org/10.1016/j.combiomed.2014.04.010>.
- [68] Zarringhalam K, Enayetallah A, Reddy P, Ziemek D. Robust clinical outcome prediction based on Bayesian analysis of transcriptional profiles and prior causal networks. *Bioinformatics* 2014;30(12):69–77. <https://doi.org/10.1093/bioinformatics/btu272>.
- [69] Constantinou AC, Freestone M, Marsh W, Fenton N, Coid J. Risk assessment and risk management of violent reoffending among prisoners. *Expert Syst Appl* 2015; 42(21):7511–29. <https://doi.org/10.1016/j.eswa.2015.05.025>.
- [70] Yin Z, Zhao Y, Lu X, Duan H. A hybrid intelligent diagnosis approach for quick screening of Alzheimer's disease based on multiple neuropsychological rating scales. *Comput Math Methods Med* 2015;2015. <https://doi.org/10.1155/2015/258761>.
- [71] Alobaidi ATS, Mahmood NT. Modified full Bayesian networks classifiers for medical diagnosis. *International Conference on Advanced Computer Science Applications and Technologies, ACSAT 2013* 2013:5–12. <https://doi.org/10.1109/ACSAT.2013.10>.
- [72] Vila-Francés J, et al. Expert system for predicting unstable angina based on Bayesian networks. *Expert Syst Appl* 2013;40(12):5004–10. <https://doi.org/10.1016/j.eswa.2013.03.029>.
- [73] Tylman W, et al. Real-time prediction of acute cardiovascular events using hardware-implemented Bayesian networks. *Comput Biol Med* 2016;69:245–53. <https://doi.org/10.1016/j.combiomed.2015.08.015>.
- [74] Ducher M, et al. Comparison of a Bayesian network with a logistic regression model to forecast IgA nephropathy. *Biomed Res Int* 2013;2013. <https://doi.org/10.1155/2013/686150>.
- [75] Fröhlich H, Bahamondez G, Götschel F, Korf U. Dynamic Bayesian network modeling of the interplay between EGFR and hedgehog signaling. *PLoS One* 2015;10(11):1–14. <https://doi.org/10.1371/journal.pone.0142646>.
- [76] Khademi M, Nedialkov NS. Probabilistic graphical models and deep belief networks for prognosis of breast cancer. *EEE 14th International Conference on Machine Learning and Applications, ICMIA 2015*:727–32. <https://doi.org/10.1109/ICMLA.2015.196>.
- [77] Liu S, et al. Quantitative analysis of breast cancer diagnosis using a probabilistic modelling approach. *Comput Biol Med* 2018;92(May):168–75. <https://doi.org/10.1016/j.combiomed.2017.11.014>. 2017.

- [78] Haddad T, Himes A, Campbell M. Fracture prediction of cardiac lead medical devices using Bayesian networks. *Reliab. Eng. Syst. Saf.* 2014;123:145–57. <https://doi.org/10.1016/j.res.2013.11.005>.
- [79] Jiang X, Xue D, Brufsky A, Khan S, Neapolitan R. A new method for predicting patient survivorship using efficient Bayesian network learning. *Cancer Inform* 2014;13:47–57. <https://doi.org/10.4137/CIN.S13053>.
- [80] Farmer N. An update and further testing of a knowledge-based diagnostic clinical decision support system for musculoskeletal disorders of the shoulder for use in a primary care setting. *J Eval Clin Pract* 2014;20:589–95. <https://doi.org/10.1111/jep.12153>.
- [81] Sa-ngamuang C, Haddawy P, Luvira V, Piyaphanee W, Iamsirithaworn S, Lawpoolsri S. Accuracy of dengue clinical diagnosis with and without NS1 antigen rapid test: comparison between human and Bayesian network model decision. *PLoS Negl Trop Dis* 2018;12(6):1–14. <https://doi.org/10.1371/journal.pntd.0006573>.
- [82] Yet B, Bastani K, Raharjo H, Livfvergren S, Marsh W, Bergman B. Decision support system for Warfarin therapy management using Bayesian networks. *Decis Support Syst* 2013;55(2):488–98. <https://doi.org/10.1016/j.dss.2012.10.007>.
- [83] Velikova M, Lucas PJF, Samulski M, Karssemeijer N. On the interplay of machine learning and background knowledge in image interpretation by Bayesian networks. *Artif Intell Med* 2013;57(1):73–86. <https://doi.org/10.1016/j.artmed.2012.12.004>.
- [84] Kalet AM, Gennari JH, Ford EC, Phillips MH. Bayesian network models for error detection in radiotherapy plans. *Phys Med Biol* 2015;60(7):2735–49. <https://doi.org/10.1088/0031-9155/60/7/2735>.
- [85] Fuster-Parra P, Tauler P, Bannasar-Veny M, Ligeza A, López-González AA, Aguiló A. Bayesian network modeling: a case study of an epidemiologic system analysis of cardiovascular risk. *Comput Methods Programs Biomed* 2016;126:128–42. <https://doi.org/10.1016/j.cmpb.2015.12.010>.
- [86] Cypko MA, et al. Validation workflow for a clinical Bayesian network model in multidisciplinary decision making in head and neck oncology treatment. *Int J Comput Assist Radiol Surg* 2017;12(11):1959–70. <https://doi.org/10.1007/s11548-017-1531-7>.
- [87] Zariakas V, Papageorgiou E, Regner P. Bayesian network construction using a fuzzy rule based approach for medical decision support. *Expert Syst.* 2015;32(3):344–69. <https://doi.org/10.1111/exsy.12089>.
- [88] Qian X, Wang Y, Jiang X. Parallel Bayesian network modelling for pervasive health monitoring system. the International Parallel and Distributed Processing Symposium, IPDPS 2014:1631–7. <https://doi.org/10.1109/IPDPSW.2014.182>.
- [89] Takenaka S, Aono H. Prediction of postoperative clinical recovery of drop foot attributable to Lumbar degenerative diseases, via a Bayesian network. *Clin Orthop Relat Res* 2017;475(3):872–80. <https://doi.org/10.1007/s11999-016-5180-x>.
- [90] Sandri M, Berchiolla P, Baldi I, Gregori D, De Blasi RA. Dynamic Bayesian networks to predict sequences of organ failures in patients admitted to ICU. *J Biomed Inform* 2014;48:106–13. <https://doi.org/10.1016/j.jbi.2013.12.008>.
- [91] Moreira LB, Namen AA. A hybrid data mining model for diagnosis of patients with clinical suspicion of dementia. *Comput Methods Programs Biomed* 2018;165:139–49. <https://doi.org/10.1016/j.cmpb.2018.08.016>.
- [92] Exarchos KP, et al. A multiscale approach for modeling atherosclerosis progression. *IEEE J. Biomed. Heal. Informatics* 2015;19(2):709–19. <https://doi.org/10.1109/JBHI.2014.2323935>.
- [93] Lee J, Lee W, Park IS, Kim HS, Lee H, Jun CH. Risk assessment for hypertension and hypertension complications incidences using a Bayesian network. *IIE Trans Healthc Syst Eng* 2016;6(4):246–59. <https://doi.org/10.1080/19488300.2016.1232767>.
- [94] Van der Heijden M, Velikova M, Lucas PJF. Learning Bayesian networks for clinical time series analysis. *J Biomed Inform* 2014;48:94–105. <https://doi.org/10.1016/j.jbi.2013.12.007>.
- [95] Ramos J, et al. NRF1 motif sequence-enriched genes involved in ER/PR –ve HER2 +ve breast cancer signaling pathways. *Breast Cancer Res Treat* 2018;172(2):469–85. <https://doi.org/10.1007/s10549-018-4905-9>.
- [96] Chang YS, Fan CT, Lo WT, Hung WC, Yuan SM. Mobile cloud-based depression diagnosis using an ontology and a Bayesian network. *Future Gener Comput Syst* 2015;43–44:87–98. <https://doi.org/10.1016/j.future.2014.05.004>.
- [97] Noyes N, Cho KC, Ravel J, Forney LJ, Abdo Z. Associations between sexual habits, menstrual hygiene practices, demographics and the vaginal microbiome as revealed by Bayesian network analysis. *PLoS One* 2018;13(1):1–25. <https://doi.org/10.1371/journal.pone.0191625>.
- [98] Lappenschaar M, et al. Multilevel temporal Bayesian networks can model longitudinal change in multimorbidity. *J Clin Epidemiol* 2013;66(12):1405–16. <https://doi.org/10.1016/j.jclinepi.2013.06.018>.
- [99] Spyroglou II, Spöck G, Rigas AG, Paraskakis EN. Evaluation of Bayesian classifiers in asthma exacerbation prediction after medication discontinuation. *BMC Res Notes* 2018;11(1):18–23. <https://doi.org/10.1186/s13104-018-3621-1>.
- [100] Aljawad DA, et al. Breast cancer surgery survivability prediction using Bayesian network and support vector machines. 2017 Int. Conf. Informatics, Heal. Technol. ICIHT 2017, no. February 2019 2017. <https://doi.org/10.1109/ICIHT.2017.7899000>.
- [101] Wang Z, et al. 3-D stent detection in intravascular OCT using a Bayesian network and graph search. *IEEE Trans Med Imaging* 2015;34(7):1549–61. <https://doi.org/10.1109/TMI.2015.2405341>.
- [102] Ducher M, Mounier-Véhier C, Lantelme P, Vaisse B, Baguet JP, Fauvel JP. Reliability of a Bayesian network to predict an elevated aldosterone-to-renin ratio. *Arch Cardiovasc Dis* 2015;108(5):293–9. <https://doi.org/10.1016/j.acvd.2014.09.011>.
- [103] Gámez-Pozo A, et al. Functional proteomics outlines the complexity of breast cancer molecular subtypes. *Sci Rep* 2017;7(1):1–13. <https://doi.org/10.1038/s41598-017-10493-w>.
- [104] Dranca L, et al. Using Kinect to classify Parkinson's disease stages related to severity of gait impairment. *BMC Bioinformatics* 2018;19(1):1–15. <https://doi.org/10.1186/s12859-018-2488-4>.
- [105] Neapolitan R, Jiang X, Ladner DP, Kaplan B. A primer on Bayesian decision analysis with an application to a kidney transplant decision. *Transplantation* 2016;100(3):489–96. <https://doi.org/10.1097/TP.0000000000001145>.
- [106] Park E, Chang HJ, Nam HS. A Bayesian network model for predicting post-stroke outcomes with available risk factors. *Front Neurol* 2018;9(September):1–11. <https://doi.org/10.3389/fneur.2018.00699>.
- [107] Cai ZQ, Guo P, Bin Si S, Geng ZM, Chen C, Cong LL. Analysis of prognostic factors for survival after surgery for gallbladder cancer based on a Bayesian network. *Sci Rep* 2017;7(1):1–10. <https://doi.org/10.1038/s41598-017-00491-3>.
- [108] Syafiandini AF, Wasito I. Metastasis identification based on clinical parameters using Bayesian network. 4th International Conference on Information and Communication Technology 2016;4:1–6. <https://doi.org/10.1109/ICoICT.2016.7571919>. no. c.
- [109] Armstrong S. The apps attempting to transfer NHS 111 online. *BMJ* 2018;360(January):1–5. <https://doi.org/10.1136/bmj.k156>.
- [110] MMC Ventures. Beyond the hype: artificial intelligence podcast Interview - Babylon health. 2017 [Online]. Available: <https://www.healthcare.digital/single-post/2018/03/29/Healing-Healthcare-with-Artificial-Intelligence-“Beyond-The-Hype”-with-Dr-Ali-Parsa-Founder-and-CEO-of-Babylon-Health#1>.
- [111] Jensen FV, Chamberlain B, Nordahl T, Jensen F. Analysis in HUGIN of data conflict. Proceedings of the Sixth Annual Conference on Uncertainty in Artificial Intelligence 1990:519–28.
- [112] Jensen K, Andreassen S. Generic probabilistic networks in medical decision support, vol. 39. IFAC; 2006.
- [113] Greenhalgh T. How to improve success of technology projects in health and social care. *Public Health Res Pract* 2018;28(2). <https://doi.org/10.17061/phrp2831815>.
- [114] Osheroff J, et al. Improving outcomes with clinical decision support: an implementer's guide. 2nd ed. Healthcare Information and Management Systems Society (HIMSS); 2012.